

CONSENT FOR ANESTHESIA AND EXTRACTION OF TEETH

Patient Name: _____ * _____ *
Last First MI Preferred Name

Planned treatment for today is * _____

Alternative Treatment methods include *

Please read each paragraph . If you have any questions, please ask your doctor BEFORE signing:

You have the right to be informed about your diagnosis and planed surgery so that you can decide whether to have the procedure or not after knowing the risk and benefits.

Removal of a tooth is a permanent procedure. Whether the procedure is easy or difficult, it is still a surgical procedure. All surgeries have some risk.

They include the following :

1. Generalized Swelling, bruising and pain at the extraction site and surrounding area.
2. Stretching of the corners of the mouth that may lead to cracking or bruising.
3. Possible infection that may need more treatment
4. Dry socket - a jaw pain beginning in a few days after surgery, needing additional care
5. Possible damage to teeth close to the ones being taken out, more often those with large fillings or crowns
6. Numbness, pain, or changed feelings in the teeth, gums, lip, chin and/or tongue (including possible loss of taste). This is due to the closeness of tooth roots (mainly Wisdom teeth) to the nerves, which can be injured or damaged. Usually the numbness goes away but in some cases can be permanent.
7. Trismus - you can only move your mouth a little. This is most common after wisdom teeth are taken out. Sometimes it happens because of jaw joint problems (TMJ), mainly when TMJ disease is already there.
8. Bleeding - oozing can often happen for several hours, but a lot of bleeding is not common.
9. Sharp ridges or bone splinters may form later at the edge of the hole where the tooth was taken out. These may need another surgery to smooth or remove.
10. Sometimes tooth roots may be left in to avoid harming important things such as nerves or a sinus (a hollow place above your upper back teeth)
11. The roots of the upper back teeth are often close to the sinus. Sometimes a piece of root can get into the sinus . An opening may occur from the sinus into the mouth that may need more treatment.
12. It is very rare that the jaw will break, but it is possible in cases where the tooth are buried very deep in their sockets.

* I have read the information above regarding Anesthesia and I give my Consen for this to be applied

Response Date: _____